

Supernova search at intermediate z

II. Host galaxy morphology

J. Méndez (1,2), P. Ruiz-Lapuente (1), G. Altavilla (1),
A. Balastegui (1), M. Irwin (3), K. Schahmanec (4), C. Balland (5,6),
R. Pain (4), N. Walton (3)

1) *Department of Astronomy, CER for Astrophysics, Particle Physics and Cosmology, University of Barcelona, Diagonal 647, E-08028, Barcelona, Spain*

2) *Isaac Newton Group of Telescopes, 38700 Santa Cruz de La Palma, Islas Canarias, Spain*

3) *Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge. CB3 0HA, United Kingdom*

4) *LPNHE-IN2P3-CNRS-Universités Paris 6 et Paris 7, 4 place Jussieu, 75252 Paris Cedex 05 France*

5) *Institut d'Astrophysique Spatiale, Bâtiment 121, Université Paris 11, 91405 Orsay Cedex, France*

6) *Université Paris Sud, IAS-CNRS, Bâtiment 121, Orsay Cedex, France*

Abstract. We discuss the host galaxy morphology of the 8 SNe discovered as a part of the International Time Programme (ITP) project “ Ω and Λ from Supernovae, and the Physics of Supernovae Explosions” at the European Northern Observatory (ENO). Identification of the SN host galaxy types was done exploiting both imaging and spectroscopy. A peculiar SNIa at $z=0.033$ is found in a spiral galaxy, as most other SNeIa with z between 0.1 and 0.4. A complete account of these studies will be given elsewhere.

1. Galaxy morphology

Spectroscopic observations were carried out using the Intermediate dispersion Spectrograph and Imaging System (ISIS) mounted at the 4.2m WHT. Imaging was done using the WFC at the 2.5m INT, ALFOSC at the 2.5m NOT and DOLORES at the 3.6m TNG. From the analysis of the galaxy spectra, though contaminated by the SN, we can discriminate between spheroidal (elliptical and lenticular) and spiral galaxies. In the analysed SN sample only the morphologies of a few host galaxies can be easily identified by visual inspection (see Fig.1, 2). The results are summarized in Table 1¹.

¹Host galaxy images of these SNeIa can be found at the web site of this project.




Figure 1. SN 2002lq host galaxy image (left panel) and spectrum (right panel). The SN position is marked by an arrow. The SN spectrum shows numerous galaxy features (Balmer series lines are clearly visible). Both spectroscopy and imaging are consistent with a face-on spiral galaxy.




Figure 2. SN 2002lk host galaxy image (left panel) and spectrum (right panel). The SN position is marked by an arrow. The SN spectrum shows a few galaxy features (due to the high brightness of the SN, the only galaxy lines clearly identified are H_α , nitrogen and sulphur lines). A deep sodium absorption is visible too. Spectroscopy and the absorption lane visible in the imaging suggest that this is an edge-on spiral galaxy.

Table 1. Host galaxy summary.

SN name	z	SN Type	Galaxy Type	Offset from nucleus	Identified lines
2002li	0.329	Ia	Spiral	0".1 W 0".2 S	H_α , H_β , H_δ , [SII], [NII], [OIII], [OII]
2002lj	0.180	Ia	Spheroidal	0".2 W	-
2002lp	0.144	Ia	Spiral	0".2 E 2".0 N	H_α , H_γ , H_θ , [OII]
2002lq	0.269	Ia	Spiral	4".5 E 0".7 S	H_α , H_β , H_γ , H_δ , H_ζ , H_η , H_θ , [SII], [OIII]
2002lr	0.255	Ia	Spiral?	3".0 S	H_β , [SII], [NII], [NeV]
2002lk	0.033	Ia pec.	Spiral	0".7 W 0".2 N	H_α , [SII], [NII], NaD
2002ln	0.138	II	Spiral	1".1 W 8".3 S	H_α , H_β , H_ζ , H_θ , [SII]
2002lo	0.136	II	Spiral	0".6 E 1".3 S	H_α , H_β , H_θ , [NII], [OIII], [OII]